



# Non-destructive estimation of leaf area and leaf weight of *Cinchona officinalis* L. (Rubiaceae) based on linear models

Annick Estefany Huaccha-Castillo, Franklin Hitler Fernandez-Zarate, Luis Jhoseph Pérez-Delgado, Karla Saith Tantalean-Osores, Segundo Primitivo Vaca-Marquina, Tito Sanchez-Santillan, Eli Morales-Rojas, Alejandro Seminario-Cunya & Lenin Quiñones-Huatangari

To cite this article: Annick Estefany Huaccha-Castillo, Franklin Hitler Fernandez-Zarate, Luis Jhoseph Pérez-Delgado, Karla Saith Tantalean-Osores, Segundo Primitivo Vaca-Marquina, Tito Sanchez-Santillan, Eli Morales-Rojas, Alejandro Seminario-Cunya & Lenin Quiñones-Huatangari (2023) Non-destructive estimation of leaf area and leaf weight of *Cinchona officinalis* L. (Rubiaceae) based on linear models, *Forest Science and Technology*, 19:1, 59-67, DOI: [10.1080/21580103.2023.2170473](https://doi.org/10.1080/21580103.2023.2170473)

To link to this article: <https://doi.org/10.1080/21580103.2023.2170473>



© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 25 Jan 2023.



[Submit your article to this journal](#)



Article views: 2018



[View related articles](#)










[View Crossmark data](#)



Citing articles: 4 [View citing articles](#)

## Non-destructive estimation of leaf area and leaf weight of *Cinchona officinalis* L. (Rubiaceae) based on linear models

Annick Estefany Huaccha-Castillo<sup>a</sup> , Franklin Hitler Fernandez-Zarate<sup>b</sup> , Luis Jhoseph Pérez-Delgado<sup>c</sup>, Karla Saith Tantalean-Osores<sup>c</sup>, Segundo Primitivo Vaca-Marquina<sup>d</sup> , Tito Sanchez-Santillan<sup>e</sup> , Eli Morales-Rojas<sup>f</sup> , Alejandro Seminario-Cunya<sup>b</sup>  and Lenin Quiñones-Huatangari<sup>a</sup> 

<sup>a</sup>Instituto de Ciencia de Datos, Universidad Nacional de Jaén, Cajamarca, Perú; <sup>b</sup>Facultad de Ciencias Agrarias, Universidad Nacional Autónoma de Chota, Cajamarca, Perú; <sup>c</sup>Escuela Profesional de Ingeniería Forestal y Ambiental, Universidad Nacional de Jaén, Cajamarca, Perú; <sup>d</sup>Facultad de Ciencias Agrarias, Universidad Nacional de Cajamarca, Cajamarca, Perú; <sup>e</sup>Instituto de investigaciones de la Amazonía Peruana, Iquitos, Perú; <sup>f</sup>Facultad de Ciencias Naturales y Aplicadas, Universidad Nacional Intercultural Fabiola Salazar Leguía de Bagua, Bagua, Perú

### ABSTRACT

Non-destructive methods that accurately estimate leaf area (*LA*) and leaf weight (*LW*) are simple and inexpensive, and represent powerful tools in the development of physiological and agronomic research. The objective of this research is to generate mathematical models for estimating the *LA* and *LW* of *Cinchona officinalis* leaves. A total of 220 leaves were collected from *C. officinalis* plants 10 months after transplantation. Each leaf was measured for length, width, weight, and leaf area. Data for 80% of leaves were used to form the training set, and data for the remaining 20% were used as the validation set. The training set was used for model fit and choice, whereas the validation set allowed assessment of the model's predictive ability. The *LA* and *LW* were modeled using seven linear regression models based on the length (*L*) and width (*Wi*) of leaves. In addition, the models were assessed based on calculation of the following statistics: goodness of fit ( $R^2$ ), root mean squared error (*RMSE*), Akaike's information criterion (*AIC*), and the deviation between the regression line of the observed versus expected values and the reference line, determined by the area between these lines (*ABL*). For *LA* estimation, the model  $LA = 11.521(Wi) - 21.422$  ( $R^2 = 0.96$ ,  $RMSE = 28.16$ ,  $AIC = 3.48$ , and  $ABL = 140.34$ ) was chosen, while for *LW* determination,  $LW = 0.2419(Wi) - 0.4936$  ( $R^2 = 0.93$ ,  $RMSE = 0.56$ ,  $AIC = 37.36$ , and  $ABL = 0.03$ ) was selected. Finally, the *LA* and *LW* of *C. officinalis* could be estimated through linear regression involving leaf width, proving to be a simple and accurate tool.

### ARTICLE HISTORY

Received 31 October 2022  
Accepted 15 January 2023

### KEYWORDS

*Cinchona* tree; leaf dimensions; ImagJ software; leaf morphology; mathematical models

## Introduction

*Cinchona officinalis* is a plant species of high medicinal value because its bark contains alkaloids such as quinine, which has been used as an effective treatment for malaria for more than 300 years (Córdor et al. 2009; Canales et al. 2020). In Peru, there are few forest relicts with *C. officinalis* located mainly in the regions of Cajamarca and Piura (Huamán 2020; Fernandez-Zarate et al. 2022). This species is being decimated by agriculture, livestock and logging (Zevallos 1989; Castañeda et al. 2019; Carbajal 2022; Fernandez et al. 2022), conditions that have led to the implementation of conservation and recovery programs in Peru (Albán-Castillo et al. 2020). The success of these programs will depend largely on the silvicultural management of the species at the nursery level, in that sense, the study of leaf area (*LA*) allows the analysis of eco-physiological factors such as transpiration, nutrient assimilation percentage, photosynthetic efficiency,

specific leaf area and leaf area index, which is reflected in crop growth (Torri et al. 2009; Keramatlou et al. 2015; Montelatto et al. 2020), while leaf weight (*LW*) will allow measuring the relative growth rate (Suárez et al. 2018) and biomass accumulation (Tieszen 1982).

Mathematical models describing the relationship between plant growth, dry matter generation, and leaf area (*LA*) are often used to study crop growth. Analysis of these variables over time allows assessing the effect of environmental and management conditions on crop development (Suárez et al. 2018). Destructive sampling is often required for studies that evaluate plant growth; however, this is impossible in trials whose purpose it is to monitor leaf growth over a given period (Falvo et al. 2008; Kishor Kumar et al. 2017; Suárez et al. 2018).

There are direct and indirect methods to determine the *LA* of plants (Kishor Kumar et al. 2017). Of the direct methods, the use of conventional planimeters, scanners, or fixed cameras and software for image

**CONTACT** Franklin Hitler Fernandez Zarate  [fran.9615fernandez@gmail.com](mailto:fran.9615fernandez@gmail.com)  Facultad de Ciencias Agrarias, Universidad Nacional Autónoma de Chota, Cajamarca, Perú.

© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

processing stand out (Granier et al. 2002; Liu et al. 2017; Suárez et al. 2018). However, these methods require leaf extraction (Cristofori et al. 2007; Fallovo et al. 2008). On the other hand, most non-destructive methods allow estimation of  $LA$  and leaf weight through regression models considering leaf length and width (Pompelli et al. 2012; Tondjo et al. 2015; Weraduwege et al. 2015; Kishor Kumar et al. 2017; Liu et al. 2017). Regression models that consider leaf structural parameters such as length ( $L$ ),  $L^2$ , width ( $Wi$ ),  $Wi^2$ , or a combination of length and width ( $L \times Wi$ ) have been found to have high accuracy in estimating  $LA$  (Cristofori et al. 2007; Meng et al. 2019). Leaf weight has commonly been determined through oven drying, a method that is destructive and time-consuming (Dwyer et al. 2014; Freschet et al. 2015).

Models for estimating  $LA$  as a function of leaf length and width have been generated, for example, in coffee (Antunes et al. 2008), kaffir lime (Budiarto et al. 2022), cocoa (Suárez et al. 2018), grape (Tsialtas et al. 2008), strawberry (Demirsoy et al. 2005), hazelnut (Cristofori et al. 2007), walnut (Zellers et al. 2012), cherry (Demirsoy et al. 2005), and other species of small fruits (Fallovo et al. 2008). However, during the literature review no reports on the subject under study were found for species of the genus *Cinchona*. Under this context, the objective of this research is to generate a mathematical model that non-destructively estimates the  $LA$  and  $LW$  of *C. officinalis* L. (Rubiaceae), taking into account leaf parameters (leaf length and width) as independent variables.

## Materials and methods

### Study area

Data on the  $L$ ,  $Wi$  and weight of leaves of *C. officinalis* were collected and processed in June 2022 from the Fernandez Zarate family forest nursery, located in the community La Cascarilla ( $5^{\circ}40'21.12''$  S and  $78^{\circ}53'55.65''$  W) province of Jaen at 1810 m, which is characterized by annual precipitation of 1730 mm, minimum temperature of  $13^{\circ}\text{C}$ , and maximum temperature of  $20.5^{\circ}\text{C}$  (Fernandez et al. 2022).

### Characteristics of the leaves of *C. officinalis*

They are elliptic-ovate, simple, opposite and decussate, with petiole, acute or acuminate apex, with entire margin (Zevallos 1989; Huamán 2020).

### Data collection

Fifty-five *C. officinalis* plants were randomly selected at nursery level and were 10 months old. Four leaves per plant were extracted (Figure 1), in total 220 leaves were collected without visible damage that could alter their shape (Suárez et al. 2018).

The fresh weight of each selected leaf was then determined using a Kmt Style electronic balance ( $200 \pm 0.01$  g). In addition, each leaf was photographed according to the methodology described in




Figure 1. Location of the two pairs of leaves sampled per plant of *C. officinalis*.





Figure 2. Diagram of *C. officinalis* leaf showing length ( $L$ ) and width ( $Wi$ ) measurement positions.

(Fernandez-Zarate et al. 2022). Leaf length ( $L$ ) from base to apex and maximum leaf width ( $Wi$ ) were measured using ImageJ software (Figure 2) (Baker et al. 1996). These measurements were independently performed by three people in order to decrease bias in image processing.

### Statistical analysis

Box and whisker plots were used for each morphometric variable evaluated ( $L$ ,  $Wi$ ,  $LA$ , and  $LW$ ). In



**Figure 3.** Descriptive analysis of the independent variables length ( $L$ ) and width ( $W$ ) (A) and dependent variables leaf area ( $LA$ ) (B) and weight ( $LW$ ) (C) of *C. officinalis* leaves.

addition, since the data were not normally distributed, pairwise Spearman correlation coefficients were calculated between the sets of independent variables ( $L$ ,  $W$ ,  $L^2$ ,  $W^2$ ,  $(L + W)^2$ , and  $L \times W$ ) and dependent variables ( $LA$  and  $LW$ ).

Linear regression was performed between the dependent variables ( $LA$ ) and  $LW$  with different independent variables, including  $L$ ,  $W$ ,  $L^2$ ,  $W^2$ ,  $(L + W)^2$ , and the product  $L \times W$  (Keramatlou et al. 2015). Finally, the root mean squared error ( $RMSE$ ) and Akaike's information criterion ( $AIC$ ) were calculated.

### Model validation

Of the 220 leaves sampled, data from 80% of the leaves were randomly selected to form a training set that was used to determine the models, and data of the remaining 20% were used as a validation set to estimate the predictive capacity of the fitted model (Suárez et al. 2022). The mathematical model was chosen taking into account the highest coefficient of determination ( $R^2$ ) and the lowest  $RMSE$  and  $AIC$ . In addition, in order to analyze trends in the deviation of observed to expected values, scatter plots of observed versus expected values were made, and a reference line  $y = x$  and the regression line of the observed versus expected values were superimposed. The deviation between the regression line of the observed versus the expected reference line indicated that there was bias. To determine this bias, the area between these lines ( $ABL$ ) was calculated; the

lower the  $ABL$ , the lower the bias, thereby indicating the higher performance of the model in terms of more accurate prediction (Suárez et al. 2018).

### Results

Figure 3 shows the descriptive analyses demonstrated by box and whisker plots of the variables that were used to determine the linear model for estimating leaf area and leaf weight of *C. officinalis* in a non-destructive way. The box plot graphs show the distribution of  $L$  and  $W$  (Figure 3(A)),  $LA$  (Figure 3(B)), and  $LW$  (Figure 3(C)) as a function of the range of quartiles and outliers. The circles correspond to the extreme values; that is, those far away from the median in the  $W$ ,  $LW$ , and  $LA$  data.

In addition, Spearman correlation coefficients (since the data are not normally distributed) were determined pairwise between the set of morphometric variables (Figure 4). The variable  $L \times W$  showed a higher correlation for estimating  $LA$  and  $LW$  ( $r_s = 1$  and  $r_s = 0.98$ , respectively).

Descriptive analysis was carried out of the independent variables of length ( $L$ ) and width ( $W$ ) and dependent variables of leaf area ( $LA$ ) and weight ( $LW$ ) of *C. officinalis* leaves.

The linear correlations between the independent variables ( $L$ ,  $W$ ,  $L^2$ ,  $W^2$ ,  $(L + W)^2$ ,  $L \times W$ ) and the dependent variable  $LA$  are shown in Figure 5(A-F).




Figure 4. Matrix of correlation diagrams using Spearman's test between LA, LW, L, and Wi of *C. officinalis* leaves. (\*\*\*) indicate a  $p < 0.001$ .





Figure 5. Linear regression models predicting the leaf area of *C. officinalis* as a function of the independent variables (L, Wi, L<sup>2</sup>, W<sup>2</sup>, (L + W)<sup>2</sup>, and L × Wi).

**Table 1.** Summary of the evaluation of the six models in predicting leaf area of *C. officinalis* based on the training data set.

Model code	Mathematical model	$R^2$	RMSE	AIC	ABL
Model 1	$LA = 6.8316(L) - 28.181$	0.925	28.512	93.327	1564.855
Model 2	$LA = 11.521(Wi) - 22.422$	0.964	28.162	3.480	140.336
Model 3	$LA = 0.3796(L^2) - 1.5169$	0.959	28.794	318.500	1884.417
Model 4	$LA = 1.1088(W^2) + 2.4448$	0.975	28.918	197.963	1604.466
Model 5	$LA = 0.1559(L+Wi)^2 - 0.7262$	0.994	29.072	317.987	1983.769
Model 6	$LA = 0.6724(L \times Wi) - 0.1422$	0.998	29.101	138.992	2128.006

**Figure 6.** Linear regression models predicting leaf weight of *C. officinalis* as a function of the independent variables ( $L$ ,  $Wi$ ,  $L^2$ ,  $Wi^2$ ,  $(L + Wi)^2$ , and  $L \times Wi$ ).

It is evident that the parameters are highly correlated with  $LA$  ( $R^2 = 0.925 - 0.998$ ). Table 1 shows the regression coefficients,  $RMSE$ , and  $AIC$  of the six models used to estimate  $LA$ . With respect to the statistics calculated to select a particular model, none of the models comply for all the considered statistics. Model 6 achieved the highest  $R^2$ , whereas Model 2 had the lowest  $RMSE$  and  $AIC$ .


The linear correlations between the independent variables ( $L$ ,  $Wi$ ,  $L^2$ ,  $Wi^2$ ,  $(L + Wi)^2$ ,  $L \times Wi$ ) and the dependent variable  $LW$  are shown in Figure 6(A-F). It is evident that the parameters are highly correlated

with  $LW$  ( $R^2 = 0.89 - 0.97$ ). Table 2 shows the regression coefficients,  $RMSE$ , and  $AIC$  of the six models used to estimate the  $LW$  of *C. officinalis* leaves. With respect to the statistics calculated to select a particular model, it can be seen that none of the models showed compliance for all the considered statistics. Model 6 had the highest  $R^2$ , Model 1 achieved the lowest  $RMSE$ , and Model 2 had the lowest  $AIC$ .

Deviation of the regression line from the expected versus observed values from the baseline is evidence of bias, as shown in Figures 7 and 8. In this case, the baseline and regression line were very close in the

**Table 2.** Summary of the evaluation of the six models in predicting leaf weight of *C. officinalis* based on the training data set.

Model code	Mathematical model	$R^2$	RMSE	AIC	ABL
Model 1	$LW = 0.1433(L) - 0.6136$	0.888	0.154	123.778	0.329
Model 2	$LW = 0.2419(Wl) - 0.4936$	0.928	0.560	37.359	0.029
Model 3	$LW = 0.0080(L^2) - 0.0563$	0.929	0.560	362.675	0.428
Model 4	$LW = 0.0234(W^2) + 0.0272$	0.944	0.563	211.452	0.609
Model 5	$LW = 0.0033(L+Wl)^2 - 0.0397$	0.963	0.567	411.447	0.423
Model 6	$LW = 0.0142(L \times Wl) - 0.0274$	0.966	0.567	271.552	0.400

**Figure 7.** Relationship between expected leaf area and observed leaf area of leaves of *C. officinalis* plants. Regression line of observed versus expected values (solid line) and dotted reference line ( $x=y$ ).


visual analysis of the differences between a set of competing models. When modeling the LA of *C. officinalis* and considering the  $R^2$ , RMSE (smaller, better), and AIC (smaller, better), two models (Model 2 and Model 6) showed some outstanding features. However, Model 2 had only one input parameter, and the regression line of the observed versus predicted was closer to the baseline (lower ABL, Figure 7). Therefore, Model 2 is proposed as the best of the six evaluated models for estimating the LA of *C. officinalis*.

When modeling the LW of *C. officinalis* and considering the  $R^2$ , RMSE (smaller, better), and AIC (smaller,

better), three models (Models 1, 2, and 6) showed some outstanding features. However, Models 1 and 2 had only one input parameter, and of these, the regression line of the observed to predicted for Model 2 was closer to the baseline (lower ABL, Figure 8). Therefore, we propose that Model 2 is the best of the six evaluated models for the estimation of *C. officinalis* LW.

## Discussion

LA is used to infer plant biomass accumulation (Weraduwege et al. 2015) in addition to estimating leaf



**Figure 8.** Relationship between expected and observed leaf weights of *C. officinalis* plants for the six evaluated models. Regression line of observed versus expected values (solid line) and dotted reference line ( $x=y$ ).

growth (Nihayati et al. 2018; Budiarto et al. 2022). The calculation of *LA* and *LW* through non-destructive methods has been used in various physiological studies (photosynthetic capacity) and to assess the agronomic behavior of plants (fertilization intensity, water availability) (Lizaso et al. 2003; Swart et al. 2004; Blanco and Folegatti 2005; Roupael et al. 2007; Suárez et al. 2018). For this reason, the *L* and *Wi* of a leaf have been used in regression as predictors of leaf variables that are more complex to measure non-destructively (*LA* and *LW*) (Ma et al. 1992; Serdar and Demirsoy 2006; Roupael et al. 2007; Bakhshandeh et al. 2011; Erdoğan 2012; Pompelli et al. 2012; Fascella et al. 2013; Nehbandani et al. 2013; Keramatlou et al. 2015; Oliveira et al. 2015; Pezzini et al. 2018; Suárez et al. 2018; Wang et al. 2019; Montelatto et al. 2020; Sabouri et al. 2022).

The research results showed a high correlation between the independent variables and dependent

variables, with Spearman correlation coefficients ranging from 0.98 to 1 for the estimation of *LA* and from 0.96 to 0.98 for the estimation of *LW* of *C. officinalis*. Similarly, Spearman correlation coefficients of 0.93 have been reported between the weight and length of a leaf and of 0.89 between the weight and width (Suárez et al. 2018), whereas Serdar and Demirsoy (2006) determined a high correlation between the width, length, and area of chestnut leaves, with correlation coefficients ranging from 0.95 to 0.98.

The highest  $R^2$  values (0.998 and 0.966) were obtained for the linear model using the product between leaf length and leaf width ( $L \times Wi$ ) to estimate the *LA* and *LW* of *C. officinalis*, respectively, however, this behavior is different when calculating the *RMSE* and *AIC*, this is evidence that the models either underestimate or overestimate the *LA* or *LW* of the leaves (Basak et al. 2019; Mela et al. 2022). In this study, in addition to  $R^2$  and *RMSE*, we calculated the



*AIC* and the area between the fitted versus predicted line and the baseline (*ABL*) as additional criteria for selecting a model, taking into account bias in the same way as reported by (Suárez et al. 2018).

Considering models that were superior in more than one statistic, either obtaining a higher  $R^2$  or a lower *RMSE*, *AIC*, or *ABL*, we found that the best model to predict *LA* and *LW* of *C. officinalis* leaves is Model 2, which used as the independent variable leaf width, since it had a lower *RMSE*, *AIC*, and *ABL* (for estimating *LA*) and lower *AIC* and *ABL* (for estimating *We*). Similar results were reported by (Sabouri et al. 2022), who found that models using the *L* or *Wi* of the leaves provided more accurate estimates of *LA* and *LW*, Cristofori et al. (2007) found  $R^2$  values between 0.70 and 0.81 when estimating the *LA* as a function of the *L* or *Wi* of apple leaves, resulting in higher  $R^2$  when *Wi* of leaves was used as independent variable, for the case of rose leaves, Roupheal et al. (2007) showed several models that allowed estimating *LA* and *LW* as a function of leaf *L* and *Wi*, Roupheal et al. (2007) developed three mathematical models that estimated *LA*, fresh and dry weight of maize leaves from measurements of leaf *L* and *Wi*, resulting in strong relationships between *L* and *Wi* with *LA* and *LW* ( $R^2 > 0.85$ ).

## Conclusion

The results obtained in this research showed that *LA* and *LW* of leaves of *C. officinalis* can be estimated through linear regression based on leaf width. Based on  $R^2$ , *RMSE*, *AIC* and *ABL*, Model 2 is recommended for estimating both the *LA* ( $LA = 11.521(Wi) - 21.422$ ) and *LW* ( $LW = 0.2419(Wi) - 0.4936$ ). The use of these models in estimation and their validation was based on a data set obtained for this purpose. The equations were shown to be a simple, accurate, and time-saving tool for evaluating the growth of *C. officinalis* plants. There are some limitations regarding accuracy; however, the use of a larger amount of data would decrease extent of deviation. In addition, to increase the model accuracy, the incorporation of environmental factors, nursery management practices, and other growth factors is suggested. Finally, future work should test the applicability of the suggested models to other species of the genus *Cinchona*.


## Disclosure statement

No potential conflict of interest was reported by the author(s).

## ORCID

Annick Estefany Huaccha-Castillo  <http://orcid.org/0000-0002-9556-0080>

Franklin Hitler Fernandez-Zarate  <http://orcid.org/0000-0002-8203-9026>

Segundo Primitivo Vaca-Marquina  <http://orcid.org/0000-0001-6276-5146>

Tito Sanchez-Santillan  <http://orcid.org/0000-0002-3352-341X>

Eli Morales-Rojas  <http://orcid.org/0000-0002-8623-3192>

Alejandro Seminario-Cunya  <http://orcid.org/0000-0002-5051-729X>

Lenin Quiñones-Huatangari  <http://orcid.org/0000-0002-0953-328X>

## References

- Albán-Castillo J, Chilquillo E, Melchor-Castro B, Arakaki M, León B, Suni M. 2020. *Cinchona* L. "Árbol de la Quina": repoblamiento y reforestación en el Perú. *Rev Peru Biol.* 27(3):423–426.
- Antunes W c, Pompelli M f, Carretero D m, DaMatta F m 2008. Allometric models for non-destructive leaf area estimation in coffee (*Coffea arabica* and *Coffea canephora*). *Ann Appl Biol.* 153(1):33–40.
- Baker B, Olszyk DM, Tingey D. 1996. Digital image analysis to estimate leaf area. *J Plant Physiol.* 148(5):530–535.
- Bakhshandeh E, Kamkar B, Tsialtas JT. 2011. Application of linear models for estimation of leaf area in soybean [*Glycine max* (L.) Merr]. *Photosynth.* 49(3):405.
- Basak JK, Qasim W, Okyere FG, Khan F, Lee YJ, Park J, Kim HT. 2019. Regression analysis to estimate morphology parameters of pepper plant in a controlled greenhouse system. *J Biosyst Eng.* 44(2):57–68.
- Blanco FF, Folegatti MV. 2005. Estimation of leaf area for greenhouse cucumber by linear measurements under salinity and grafting. *Sci Agric (Piracicaba, Braz.).* 62(4):305–309.
- Budiarto R, Poerwanto R, Santosa E, Efendi D, Augusta A. 2022. Comparative and correlation analysis of young and mature kaffir lime (*Citrus hystrix* DC). *IJPB.* 13(3):270–280.
- Canales NAG, Hansen TN, Cornett C, Walker K, Driver F, Antonelli A, Maldonado C, Nesbitt M, Barnes CJ, Rønsted N. 2020. Historical chemical annotations of *Cinchona* bark collections are comparable to results from current day high-pressure liquid chromatography technologies. *J Ethnopharmacol.* 249(112375):112375.
- Carbajal JYR. 2022. La extinción del árbol de la quina por la actividad agrícola a propósito de la tradición «Los polvos de la condesa». *El Palma Juv.* 4(4):161–175.
- Castañeda JHV, Sabaleta EL, Hernández MKYB, Meza SNV, Sánchez LMQ. 2019. Comparación de sustratos en la propagación sexual y asexual del árbol de la quina (*Cinchona officinalis*). *Rev Investig Agroproducción Sustentable.* 2(3):77–85.
- Cóndor E, de Oliveira BHL, Ochoa KR, Pinedo V. 2009. Estudio químico de los tallos de *Cinchona pubescens* Vahl. *Rev Soc Quím Perú.* 75(1):54–63.
- Cristofori V, Roupheal Y, Gyves EM, Bignami C. 2007. A simple model for estimating leaf area of hazelnut from linear measurements. *Sci Hortic.* 113(2):221–225.
- Demirsoy H, Demirsoy L, Öztürk A. 2005. Improved model for the non-destructive estimation of strawberry leaf area. *Fruits.* 60(1):69–73.
- Dwyer JM, Hobbs RJ, Mayfield MM. 2014. Specific leaf area responses to environmental gradients through space and time. *Ecology.* 95(2):399–410.
- Erdoğan C. 2012. A leaf area estimation model for faba bean (*Vicia faba* L.) grown in the Mediterranean type of climate. *Süleyman Demirel Univ J Agric Turk.* 7(1):58–63.
- Falovo C, Cristofori V, de-Gyves EM, Rivera CM, Rea R, Fanasca S, Bignami C, Sassine Y, Roupheal Y. 2008. Leaf area estimation model for small fruits from linear measurements. *horts.* 43(7):2263–2267.
- Fascella G, Darwich S, Roupheal Y. 2013. Validation of a leaf area prediction model proposed for rose. *Chilean J Agric Res.* 73(1):73–76.
- Fernandez FH, Huaccha AE, Barturén LM, Quiñones L, Sánchez T. 2022. Efecto del sustrato en la propagación sexual de *Cinchona officinalis* L. (Rubiaceae). *Ecosistemas.* 31(1): 2314–2314.
- Fernandez-Zarate FH, Huaccha-Castillo AE, Quiñones-Huatangari L, Vaca-Marquina SP, Sanchez-Santillan T, Morales-Rojas E, Seminario-Cunya A, Guelac-Santillan M,

- Barturén-Vega LM, Coronel-Bustamante D. 2022. Effect of arbuscular mycorrhiza on germination and initial growth of *Cinchona officinalis* L. (Rubiaceae). *For Sci Technol.* 18(4): 182–189.
- Freschet GT, Swart EM, Cornelissen JHC. 2015. Integrated plant phenotypic responses to contrasting above- and below-ground resources: key roles of specific leaf area and root mass fraction. *New Phytol.* 206(4):1247–1260.
- Granier C, Massonnet C, Turc O, Muller B, Chenu K, Tardieu F. 2002. Individual leaf development in *Arabidopsis thaliana*: a stable thermal-time-based programme. *Ann Bot.* 89(5):595–604.
- Huamán LK. 2020. [Evaluación morfológica y del estado de conservación de seis especies del género *Cinchona* L. (Rubiaceae) en los Andes del norte y centro de Perú ]. [Master's thesis]. Lima (Peru): Universidad Nacional Mayor de San Marcos. [accessed 2022 Sep 16]. <https://cybertesis.unmsm.edu.pe/handle/20.500.12672/15618>.
- Keramatlou I, Sharifani M, Sabouri H, Alizadeh M, Kamkar B. 2015. A simple linear model for leaf area estimation in Persian walnut (*Juglans regia* L.). *Sci Hortic.* 184:36–39.
- Kumar MS, Kumar R, Sankar V, Sakthivel T, Karunakaran G, Tripathi P. 2017. Non-destructive estimation of leaf area of durian (*Durio zibethinus*) – an artificial neural network approach. *Sci Hortic.* 219:319–325.
- Liu Z, Zhu Y, Li F, Jin G. 2017. Non-destructively predicting leaf area, leaf mass and specific leaf area based on a linear mixed-effect model for broadleaf species. *Ecol Indic.* 78:340–350.
- Lizaso JJ, Batchelor WD, Westgate ME. 2003. A leaf area model to simulate cultivar-specific expansion and senescence of maize leaves. *Field Crops Res.* 80(1):1–17.
- Ma L, Gardner FP, Selamat A. 1992. Estimation of leaf area from leaf and total mass measurements in peanut. *Crop Sci.* 32(2):467–471. [cropsci1992.0011183X003200020036x](https://doi.org/10.2307/3866666).
- Mela D, Dias MG, da ST, Ribeiro J, Martinez ACP, Zuin A. 2022. Estimation of *Thunbergia grandiflora* leaf area from allometric models. *Com Sci.* 13:e3722.
- Meng Y-Y, Shao S-C, Liu S-J, Gao J-Y. 2019. Do the fungi associated with roots of adult plants support seed germination? A case study on *Dendrobium exile* (Orchidaceae). *Glob Ecol Conserv.* 17:e00582.
- Montelatto MB, Villamagua-Vergara GC, Castanho FP, Kawakami B, Zerbinato B, Silva MA, Guerra SPS. 2020. Models for leaf area estimation of three forest species in a short coppice rotation. *Acta Ecol Sin.* 40(4):263–267.
- Nehbandani A, Soltani A, Zeinali E, Raeisi S, Najafi A. 2013. Allometric relationships between leaf area and vegetative characteristics in soybean. *Int J Agric Crop Sci IJACS.* 6(16): 1127–1136.
- Nihayati E, Rosida A, Azizah N. 2018. Growth and yield of Temulawak (*Curcuma xanthorrhiza* Roxb.) and Corn (*Zea mays* L.) with various intercropping patterns. *IJPB.* 9(1):7099.
- Oliveira RLL, Moreira AR, Costa AVA, Souza LC, Lima LGS, Silva RTL. 2015. Modelos de Determinação Não Destrutiva de Área Foliar de Feijão Caupi *Vigna unguiculata* (L.). *GSTY.* 8(2):17–27.
- Pezzini RV, Cargnelutti Filho A, Alves BM, Follmann DN, Kleinpaul JA, Wartha CA, Silveira DL. 2018. Models for leaf area estimation in dwarf pigeon pea by leaf dimensions. *Bragantia.* 77(2):221–229.
- Pompelli MF, Antunes WC, Ferreira DTRG, Cavalcante PGS, Wanderley-Filho HCL, Endres L. 2012. Allometric models for non-destructive leaf area estimation of *Jatropha curcas*. *Biomass Bioenergy.* 36:77–85.
- Rouphael Y, Colla G, Fanasca S, Karam F. 2007. Leaf area estimation of sunflower leaves from simple linear measurements. *Photosynth.* 45(2):306–308.
- Sabouri A, Bakhshipour A, Poornoori M, Abouzari A. 2022. Application of image processing and soft computing strategies for non-destructive estimation of plum leaf area. *PLoS One.* 17(7):e0271201.
- Serdar Ü, Demirsoy H. 2006. Non-destructive leaf area estimation in chestnut. *Sci Hortic.* 108(2):227–230.
- Suárez JC, Casanoves F, Di Rienzo J. 2022. Non-destructive estimation of the leaf weight and leaf area in common bean. *Agronomy.* 12(3):711.
- Suárez JC, Melgarejo LM, Durán EH, Di Rienzo JA, Casanoves F. 2018. Non-destructive estimation of the leaf weight and leaf area in cacao (*Theobroma cacao* L.). *Sci Hortic.* 229:19–24.
- Swart EAMD, Groenwold R, Kanne HJ, Stam P, Marcelis LFM, Voorrips RE. 2004. Non-destructive estimation of leaf area for different plant ages and accessions of *Capsicum annum* L. *J Hortic Sci Biotechnol.* 79(5):764–770.
- Tieszen LL. 1982. Biomass accumulation and primary production. In: Hall CJ, editors. *Tech bio-product photosynth.* United Kingdom: Pergamon Press, Oxford.
- Tondjo K, Brancheriau L, Sabatier S-A, Kokutse AD, Akossou A, Kokou K, Fourcaud T. 2015. Non-destructive measurement of leaf area and dry biomass in *Tectona grandis*. *Trees.* 29(5): 1625–1631.
- Torri SI, Descalzi C, Frusso E. 2009. Estimation of leaf area in pecan cultivars (*Carya illinoensis*). *Cienc E Investig Agrar.* 36(1):53–58.
- Tsialtas JT, Koundouras S, Zioziou E. 2008. Leaf area estimation by simple measurements and evaluation of leaf area prediction models in Cabernet-Sauvignon grapevine leaves. *Photosynth.* 46(3):452–456.
- Wang Y, Jin G, Shi B, Liu Z. 2019. Empirical models for measuring the leaf area and leaf mass across growing periods in broadleaf species with two life histories. *Ecol Indic.* 102:289–301.
- Weraduwaage SM, Chen J, Anozie FC, Morales A, Weise SE, Sharkey TD. 2015. The relationship between leaf area growth and biomass accumulation in *Arabidopsis thaliana*. *Front Plant Sci.* 6:167. [accessed 2022 Oct 4].
- Zellers CE, Saunders MR, Morrissey RC, Shields JM, Bailey BG, Dyer J, Cook J. 2012. Development of allometric leaf area models for intensively managed black walnut (*Juglans nigra* L.). *Ann for Sci.* 69(8):907–913.
- Zevallos PA. 1989. *Taxonomía, distribución geográfica y status del género Cinchona en el Perú.* 1st ed. Lima: Centro de Datos para la Conservación, Universidad Nacional Agraria La Molina.